

The election of edible and non edible crop for biodiesel feedstock in Indonesia with AHP-BCR and GC analysis

Hakun Wirawasista Aparamarta ^{a,*}, Safrina Hapsari ^a, Setiyo Gunawan ^a, Rayhan Ibnu Shiena ^a, Adya Giwangkara Ariandi ^a, Yi-Hsu Ju ^b

^a Department of Chemical Engineering, Faculty of Industrial Engineering, Institut Teknologi Sepuluh Nopember (ITS), Kampus ITS Sukolilo, Surabaya 60111, Indonesia


^b Department of Chemical Engineering National Taiwan University of Science and Technology, Taipei 10607, Taiwan

* Corresponding author: hakun2397@gmail.com

Article history

Received 24 March 2019
 Revised 25 April 2019
 Accepted 1 August 2019
 Published Online 1 October 2019

Graphical abstract



Abstract

The increasing trend of domestic fuel consumption that is not followed by domestic fuel production creating problem for Indonesian government to fulfill the shortage of fuel consumption. The high dependence of Indonesian government on imported fuel creates some problem due to the large subsidy that must be given as a result of the increasing oil prices. This makes the development of biofuel important, especially biodiesel with an abundant raw material in Indonesia. In this work, the best alternative and the composition of raw materials for making biodiesel were investigated by using Gas Chromatographic (GC) and Analytic Hierarchy Process (AHP) - benefit-cost ratio (BCR). With GC analysis, it is expected to know the content of triglyceride (TG) and free fatty acid (FFA) content in crude oil for each alternative raw material. While the AHP-BCR analysis is expected to know the best alternative to raw materials for biodiesel production. The alternative raw materials that are selected namely *Calophyllum inophyllum* and *Jatropha curcas*. The selection of alternative raw material based on oil content, yield, and raw material cost. The criteria used for benefit hierarchy structure are economic, social, raw material availability, environment and technical. As for the hierarchy of cost structure, the criteria are the price of raw materials, opportunity cost, processing cost, environmental cost, and social cost. The present study clearly proved that *C. inophyllum* and *Jatropha curcas* oil potentially become the best alternative material for biodiesel production. The result shown from benefit-cost ratio were almost identical (*Jatropha curcas*, 1.01 and *C. Inophyllum*, 1). From GC analysis, *C. Inophyllum* became the potential alternative material because it has the biggest oil content for crude (75.99%) and after purification (94.24%)

Keywords: AHP, biodiesel, benefit-cost ratio, GC analysis, TLC analysis

© 2019 Penerbit UTM Press. All rights reserved

INTRODUCTION

Increasing population, expanding urbanization causes the increasing of food and energy demand in the world. This increasing demand of food and energy is not followed by the the reserves of fossil fuel that have a declining trend. The other problem is the uses of fossil oil creating the environmental concerns. So, it is very important to develop the future energy to fulfill the demand of energy and environmental friendly. One of the future energy is biodiesel. In recent studies, raw materials of biodiesel are edible oil such as palm oil, sunflower oil, and rapeseed. The utilization of edible oil to fulfill energy demand induces problems like the sustainability of the implementation due to its competition with land, water and food (Kansedo et al., 2009; Santacesaria et al., 2012). The other reasons, it also could cause food shortage especially in growing countries and environmental problems due to the use of available arable land to produce oil crops (Atabani et al., 2013).

Recently, the use of vegetable oils falls into two applications namely food industry, and industrial uses, including biodiesel. FAO (2013) reported that the use of edible oil is mostly for food, and biodiesel markets are far behind. Thus, the rapidly growing demand for edible oils has been triggered by the food market rather than industrial

or biodiesel market. In the food industry, edible oil is used for human consumption, such as baking, margarine, and frying fats or shortening (Alston et al., 2002). Using non-edible oil for biodiesel has one major weakness. The yields and quality of biodiesel produced are less than edible oil. Both edible and non-edible oil has its advantages and disadvantages. Thus, the choosing of the most appropriate source amongst them is very important based on the optimal benefits.

In such condition, multi-criterion decision making (MCDM) methods are widely used. One of the method is The Analytic Hierarchy Process (AHP) that was introduced by Saaty (1986). This method is an effective tool for dealing with such complex decision-making problems and could endorse the decision maker to set priorities and make the greatest decision. The AHP helps to capture both subjective and objective aspects of a decision by sinking complex decisions to a series of pairwise comparisons and then synthesizing the results. The essence of AHP is that it simplifies a complex decision by decomposing the problem into a hierarchy of 'criteria' or sub-problems to be analyzed individually. One of the criteria observed is technical. This criteria was used a gases chromatography method to measure triglyceride (TG) and free fatty acid (FFA) contents in the feedstocks. TG and FFA compounds were a key parameter to make biodiesel production (refined vegetable oil) from many feedstock (aparamarta et al., 2016).

The purpose of this research is to select and prioritize the best alternative to raw materials for biodiesel production. The alternative raw materials that are selected namely *Calophyllum inophyllum* and *Jatropha curcas*. The selection of alternative raw materials was based on the oil content, yield and feedstock price (Gui, et al, 2008). They became the important factor to resolve the suitability of a crop as the raw material for biodiesel production. The reason is oil crops with higher oil yield could decrease the production cost. Generally, the cost of raw materials achieve about 70–80% of the total production cost of biodiesel (Gui et al., 2008). The criteria used for benefit hierarchy structure are economic, social, raw material availability, environment and technical. As for the hierarchy of cost structure is the price of raw materials, opportunity cost, processing cost, environmental cost, and social cost. This criteria of hierarchy benefit and cost were based on the improvement of the previous works. Ilham and Nime (2019) was evaluated the best priority of biodiesel feedstock with FFA content, oil yield, and plant availability as the criterias of AHP methods. The other criterias are government policy, plant productivity and surplus production of raw material (Hambali et al., 2016).

EXPERIMENTAL

Materials

C. inophyllum and *Jatropha curcas* oil was bought from Koperasi Jarak Tani Lestari (Klaten, Indonesia) and some forest (Wonosobo, Central Java and Cirebon, West Java, Indonesia). Thin-layer chromatography (TLC) aluminum plates were obtained from Merck (Darmstadt, Germany). Standard triolein, tripalmitin and fatty acids were purchased from Sigma Chemical Company (St. Louis, MO). All reagents and solvents were either analytical-grade or HPLC-grade were obtained from commercial sources.

Extraction of nonpolar lipids fraction (NPLF) from crude *C. inophyllum* and *Jatropha curcas* oils (Single-stage solvent extraction)

The single stage extraction was explained by Aparamarta et al., (2016). Nonpolar solvents (petroleum ether) and polar solvent (methanol) were applied in this work. The ratio of mixture solvent (non polar and polar solvent) of 3:1 was used in this work. The result of this extraction have 2 layers namely non polar lipid fraction (NPLF) and polar lipid fraction (PLF). NPLF, such as hydrocarbons and TG, were extracted by nonpolar solvents in the upper layer. The extract was separated and concentrated to give the NPLF. The polar extract was designated as the polar lipids fraction (PLF). The NPLF and PLF were then analyzed using high temperature gas chromatography (HT-GC) and thin layer chromatography (TLC).

Multistage solvent extraction (Batchwise eight-stage extraction)

Multistage solvent extraction with 8 stages was described by aparamarta et al., (2018). Stirred batchwise method was used in this work. The result of NPLF fraction in the earlier step was extracted using polar solvent (methanol) at room temperature. The adding of polar solvent in the NPLF fraction was done in eight stages to get a higher content of TG. The polar solvent extracts (1–8 stage fractions) were collected and concentrated to give the PLF. Afterward, the NPLF and PLF were analyzed using HT-GC and TLC.

Thin layer chromatography (TLC) analysis

TLC as qualitative analysis was applied to the sample as defined by Gunawan et al. (2008). TLC paper that has been colored by the sample was immersed in a mobile phase of hexane: ethyl acetate: acetic acid at 90: 10: 1 (v/v/v).

High temperature gas chromatography (HT-GC) analysis

Components in *C. inophyllum* seed oil, such as FFA, TG, and other compounds were identified using pure standard solution as defined by Aparamarta et al. (2018). Quantitative analysis of the sample was calculated on a Shimadzu GC-2010 gas chromatograph (Kyoto, Japan), equipped with a flame ionization detector. Separations were carried out

on a DB-5HT (5%-phenyl) methylpolysiloxane nonpolar column (Phenomenex, USA).

Analytic hierarchy process-benefit cost ratio (AHP-BCR) analysis

Crude of alternative raw material was extracted and tested with gas chromatography (GC) to measure the content of TG and FFA in the crude *C. Inophyllum* oil. The result of this analysis is used as primary data to be input into economic and technical criteria in benefit hierarchy of AHP analysis.

In this work, Expert Choice 11 was applied to get the finest goal from decision-making problems. It creates a model that can help the human frame which have logic, experience, knowledge and sense factors to optimized in a systematic process (Shean, 2012). This model has 3 layers; the first layer is criteria. It has 4 criteria namely economy, social, raw material availability, environmental and technical for hierarchy benefit and raw material cost, opportunity cost, operating cost, social cost and environmental cost of hierarchy cost. The second layer is sub-criteria. It has machine efficiency, material saving, energy saving, selling power, pro-growth, pro job, pro-poor, reserve, productivity, location, reduced emission, the impact for human health, increased TG and decreased FFA for hierarchy benefit. For hierarchy cost, it has two sub-criteria namely purchased cost and transportation cost

RESULTS AND DISCUSSION

Raw material is key point to implement biodiesel production in Indonesia. Generally, the cost of raw materials reach about 70–80% of the total biodiesel production cost (Kusdiana and Saka, 2002; Demirbas, 2003; Du et al, 2004; Gui et al., 2008). For this reason, yield and oil content become the important factors to decide the suitability of a crop as a raw material for biodiesel production. These factors is the key to reduce the biodiesel production cost (Gui et al., 2008). The biggest component in oil content is TG (Aparamarta et al., 2016). This compound is the key component to make a higher yield in the biodiesel production. The other key component is FFA. This compound composition in the oil should be below than 3% for alkaline-catalyzed transesterification (Ribeiro et al., 2011) The source of edible and non-edible oil can be seen on table I.

Table 1 The non edible and edible source for biodiesel feedstock.

Feedstock	Source	Oil Content (Wt%)	Yield (Kg oil/ha)	Cetane Number	Reference
Non edible oil					
<i>C. inophyllum</i>	Seed	40-73	4680	57	Singh, 2010
<i>Croton tiglium</i>	Seed	30-45	-	-	Azam et al, 2005 Berti & Johnson, 2005
<i>Cuphea</i>	Seed	20-38	900	-	Umerie et a, 2010
<i>Guizotia abyssinica</i> L. (niger)	Seed	50-60	200-300	-	Sahoo et al, 2007
<i>Jatropha curcas</i> L.	Seed	20-60	1590	33.7-51	Azam et al, 2005
<i>Madhuca indica</i>	Seed	35-50	-	-	Jean, 2011
<i>Melia azedarach</i>	Seed	10-45	-	-	Langstraat, 1976;
Edible oil					
Soybean	Seed	18-20	375	-	Pryde,1980; Hoffmann, 1989
Palm	Kernel	42	5000	-	Atasie and Akinhannie, 2009
Rapeseed	Seed	41-46	1000	-	Cakmakli and Unal, 1988

Table 1 shows the source of promising edible and non-edible oil. It can be seen on non edible sources, *Jatropha curcas* and *Calophyllum inophyllum* give the highest oil yields with 1590 and 4680 kg oil per hectare respectively. Therefore, *Jatropha curcas* and *Calophyllum inophyllum* have become a promising feedstock to fulfill energy demand. For edible sources, palm oil gives the highest oil yield with 5000 kg oil per hectare, considerably higher than other oils which are only in the range of hundreds to 1000 kg oil per hectare. Therefore, palm oil has become a promising feedstock to fulfill energy demand. *C. inophyllum* and *Jatropha curcas* have one big advantage over palm in that they were nonedible and will not diminish food supply in growing countries (Atabani et al., 2013).

Batchwise solvent extraction (BSE) method was applied in this work. BSE method was used to replace the step in the chemical process of the refined of vegetable oils production, such as degumming, neutralization, and bleaching steps. From the batchwise solvent extraction result, it was separated into two layer namely NPLF and PLF (Aparamarta, 2018). Furthermore, the content of TG and FFA were chosen as the indicator for the effective separation of NPLF from crude *C. inophyllum* seed oil. Crude alternative raw material before and after purification were tested by HT-GC and TLC to see the content and separation of TG and FFA in the crude *C. inophyllum* and *Jatropha curcas* oil. TLC and HT-GC analysis of crude oil, NPLF, and PLF are shown in Figure 1a and 1b. From this figure, it was shown that TG was successfully separated in the NPLF as confirmed by TLC and HT-GC analyses.




Fig. 1a Result of the (a) HTGC and (b) TLC developed in mobile phase (90:10:1) in the *C.inophyllum* oil




Fig. 1b Result of the (a) HTGC and (b) TLC developed in mobile phase (90:10:1) in the *Jatropha curcas* oil.

The result of this analysis is used as primary data to be input into economic and technical criteria in hierarchy benefit. With GC analysis, it obtained the TG content in *C. inophyllum* and *jatropha curcas* seed as shown in table 2. From this table, it can be seen that the TG content in the crude of *C. Inophyllum* is better than *Jatropha curcas*. TG content after purification also shows a better result on *C. Inophyllum*. The lipid content of *C. inophyllum* seeds reported in this work agrees with the earlier work. Aparamarta et al., (2017) reported that lipids content of *C. inophyllum* seeds was 70.4% .

Table 2 TG content of *C. Inophyllum* and *Jatropha curcas*' seeds.

No	Resources	TG Content (%)	
		Crude	After Purification
1	<i>Calophyllum inophyllum</i>	75.99	94.24
2	<i>Jatropha Curcas</i>	73.02	93.83

For hierarchy benefit, five types of multifaceted criteria was used namely economic, social, raw material availability, environment and technical for hierarchy benefit. Five criteria also chosen for hierarchy cost namely the price of raw materials, opportunity cost, processing cost, environmental cost, and social cost. Result of the analysis have been evaluated to determine the best alternative resources for biodiesel production in Indonesian perspective. AHP method was applied in this work to achieve the goal. This method require a procedure for the aggregation of expert opinion using the five selected criteria that are appropriate for biodiesel production. Experts involved in the assessment found that the raw material availability for hierarchy benefit and raw material cost for hierarchy cost are the most important criteria having the priority of 0.387 and 0.419, respectively. This result followed by the priorities of economic, technical, environmental and social as 0.233, 0.215, 0.127, and 0.094, respectively for hierarchy benefit and for hierarchy cost, followed by the priorities of opportunity cost, processing cost, environmental cost and social cost as 0.29, 0.166, 0.077, and 0.049, respectively. This result was agreed with previous works that surplus production of raw material became the highest criteria for biodiesel feedstocks (Hambali et al. 2015)




Fig. 2 Model structure of hierarchy benefit.




Fig. 3 Model structure of hierarchy cost.

Calophyllum inophyllum is the high amount of *oleic acid* content in *Calophyllum inophyllum* (Aparamarta et al.,2018), so its valuable to convert became biodiesel. This content was the biggest parts in the TG (Aparamarta et al.,2018). Kusmiyati and Sugiharto (2010) convert *oleic acid* to became biodiesel with a good quality to be a diesel substitute.

Table 3 AHP result for the election of raw material for biodiesel feedstock.

No	Resources	Hirarchy	Priority	Benefit-Cost Ratio
1	<i>Calophyllum inophyllum</i>	Benefit	0.699	1.05
		Cost	0.667	
2	<i>Jatropha Curcas</i>	Benefit	0.301	0.904
		Cost	0.333	

CONCLUSION

Calophyllum inophyllum became a potential resource for biodiesel production in Indonesia. It was found from GC analysis and AHP-BCR, *Calophyllum Inophyllum* has bigger TG content (94.24%) and priority (1.05) than *Jatropha Curcas*. From AHP-BCR analysis, it was found that the most important criterion for the selection of alternative resources on hierarchy benefit is raw material availability (0.387). Most important criterion for hierarchy cost is raw material cost (0.419).

ACKNOWLEDGEMENT

This work was provided by the Institute of Research and Public Services (LPPM), Institut Teknologi Sepuluh Nopember (ITS), Surabaya, Indonesia. The authors thank Mr Gunawan for valuable technical support.

REFERENCES

Aparamarta, H.W., Saputra, T., Claratika, A., Ju, Y.H., Gunawan, S. 2016. Separation and Purification of Triacylglycerols from Nyamplung (*Calophyllum inophyllum*) Oil by Batchwise Solvent Extraction. *Ind. Eng. Chem. Re.* 55, 3113-3119.

Aparamarta, H.W., Hapsari, S., Ismawan, R., Anggraeni, V., Widjaja, A., Widjaja, T., Ju, Y.H., Gunawan, S. 2018. Separation of Xanthone and Vitamin E from *Calophyllum inophyllum* Leaf. *MJFAS.* 14, 4.

Aparamarta, H.W., Angraini, D., Istianingsih, D., Susanto, D.F., Widjaja, A., Ju, Y.H., Gunawan, S. 2017. Fatty acid fragmentation of triacylglycerol isolated from crude nyamplung oil. *AIP Conference Proceedings* 1840, 060004 (2017); doi: 10.1063/1.4982284

Aparamarta, H.W., Qadariah, L., Ju, Y.H., Gunawan, S. 2018. Separation and Identification of Fatty Acid in Triacylglycerol Isolated from *Calophyllum Inophyllum* Oil. *JEAS.* 13, 2.

Alston, J.M., Chalfant, J.A., Piggott, N.E. 2002. Estimating and testing the compensated double-log demand model. *Appl. Econ.* 34, 1177-1186.

Azam, M.M., Waris, A.,Nahar, N.M. 2005. Prospects and potential of fatty acid methyl esters of some non-traditional seed oils for use as biodiesel in India. *Biomass Bioenerg.*, 29, 293-302.

Atasie, V.N., Akinhanmi, T.F., 2009. Extraction, compositional studies and physico chemical characteristics of palm kernel oil. *Pak. J. Nutr.*, 8, 800-803.

Atabani, A.E., Cesar, A.D. 2014. *Calophyllum Inophyllum* L. – A prospective non-edible biodiesel feedstock. *Studi of Biodiesel production, properties, fatty acid composition, blending and engine performance. Renew. Sust. Energ Rev.* 37, 644-655.

Berti, M.T., Johnson, B.L. 2008. Physiological changes during seed development of cuphea. *Field Crops Res.* 106, 163-170.

Boardman, A.E., Greenberg, D.H., Vining, A.R. Welmer, D.L. 2011. *Cost Benefit Analysis: Concept and practice* 4th edition.

C a k m a k li, U., Unal, M.K. 1988. Composition and mineral distribution of rapeseed varieties tested for adaptation in turkey. *Fat. Sci. Technol.*, 10, 386-389.

Gunawan, S., Fabian, C., Ju, Y.H. 2008. Isolation and purification of fatty acid steryl esters from soybean oil deodorizer distillate. *Ind. Eng. Chem. Res.* 47, 7013-7018.

Gunawan, S., Aparamarta, H.W., Kuswandi, K., Widjaja, A., Ju, Y.H. The utilization of *Xylocarpus moluccensis* seed oil as biodiesel feedstock in Indonesia. *Ind, Crops Prod.* 2013, 52, 286-291.




Fig. 4 The goal result of hierarchy benefit.




Fig. 5 The goal result of hierarchy cost

From Figure 4 and 5, It can be seen that *Calophyllum inophyllum* has a bigger priority than the *jatropha curcas* for biodiesel production with the priority of 0.699 for hierarchy benefit and 0.667 for hierarchy cost. Followed by the priority of *jatropha curcas* as 0.301 and 0.333, respectively. The consistency ratio is 0.06 for hierarchy benefit and 0.09 for hierarchy cost, so this data is acceptable (Saaty, 1990 and Saaty,1994-a).

From AHP result, it can be obtain the ratio of benefit and cost. This analyses associated with the acceptance of raw material alternative for biodiesel feedstocks. The alternative will be accept if the the total benefits should be more than the total costs incurred (Boardman et al., 2011). From Benefit cost ratio analysis, it can be seen that *Calophyllum Inophyllum* is potentially became the best alternative material for biodiesel production as can be seen on Table 3. The ratio of benefit and cost were calculated in the *Calophyllum Inophyllum* with ratio of 1.05 and in the *Jatropha Curcas* with ratio of 0.904. This work agreed with previous that the weights of *calophyllum inophyllum* is better than *Jatropha curcas* (Hambali et al.,2016). The other potential of

- Gui, M.M., Lee, K.T., Bhatia, S. 2008. Feasibility of edible oil vs.non-edible oil vs.waste edible oil as biodiesel feedstock. *Energy*. 33, 1646–53.
- Hambali, E., Nisya, F.N., Thahar, A., Nuryanti, A., Wijaya, H. 2015. Potential of Biomass as Bioenergy Feedstock in Indonesia. *Journal of the Japan Institute of Energy*. 95, 629-638.
- Hoffmann, G. 1989. The chemistry and technology of edible oils and fats and their fat products. Academic, New York, pp139-200.
- Ilham, Z., Nimme, F.H. 2019. Quantitative priority estimation model for evaluation of various nonedible plant oils as potential biodiesel feedstock. *AIMS Agriculture and Food*. 4(2): 303–319.
- Jean M. 2009. Trees in sustainable farming Auroville, TamilNadu [cited 4 February 2011]; Availablefrom: /http:// www.csfbg.org /reports / Trees%20in %20 sustainable %20 farming%20Report.pdfS.
- Kansedo, J., Lee, K.T., Bhatia, S., 2009. Cerbera odollam (sea mango) oil as a promising non-edible feedstock for biodiesel production. *Fuel*. 88, 1148-1150
- Kusmiyati, Sugiharto, A. 2010. Production of Biodiesel from Oleic Acid and Methanol by Reactive Distillation. *BCREC*, 5 , 1 – 6.
- Langstraat, A. 1976. Characteristics and composition of oil bearing materials. *J. Am. Oil Chem. Soc.* 53, 241-47.
- Pryde, E.H. 1980. Composition of soybean oil, in *Handbook of Soy Oil Processing and Utilization*, AOCS Press, Champaign, IL, pp. 13–31.
- Ribeiro, A., Castro, F., Carvalho, J. 2011. Influence of free fatty acid content in biodisel production on non edible oils In *Proceeding: WASTES: Solutions, Treatments and Opportunities*; European Environment Agency: Portugal.
- Saaty, T.L. 1986. Absolute and relative measurement with the AHP. *The most livable cities in the United States. Socio-econ. Plum Sci.* 20(6), 327-331.
- Sahoo, P.K., Das, L.M., Babu, M.K.G., Naik, S.N. 2007. Biodiesel development from high acid value polanga seed oil and performance evaluation in a CI engine. *Fuel*. 86, 448–54.
- Santacesaria, E., Vicente, G.M, Di Serio, M., Tesser. 2012. Main technologies in biodiesel production: state of the art and future challenges. *Catal. Today*.195, 2-13.
- Umerie, S.C., Okonkwo, I.F., Nwadiolor, N.A., Okonkwo, J.C. 2010. Studies on the oil and nutritive value of seeds of *Crotalaria retusa* L. (Fabaceae). *Pakistan J. Nutr.* 9, 912–4.